

HAVE A UTI?
THERE'S
BLUJEPÀ

Indication

What Is BLUJEPÀ?

BLUJEPÀ is an antibiotic used to treat women and girls 12 years of age and older weighing at least 88 pounds who have an infection of the bladder (known as an uncomplicated urinary tract infection [uUTI]), caused by certain types of bacteria.

It is not known if BLUJEPÀ is safe and effective in children under 12 years of age or weighing less than 88 pounds.

Important Safety Information

What is the most important information I should know about BLUJEPÀ?

BLUJEPÀ, an antibiotic, can cause serious side effects, including:

- **changes in electrical activity of your heart called QTc prolongation.** BLUJEPÀ can cause QTc prolongation (also called prolonged QTc interval) that can lead to serious heart rhythm problems, including Torsades de Pointes. There may be changes in your heartbeat (a fast or irregular heartbeat).


Please see Important Safety Information throughout and on pages 6 and 7.

Please see the accompanying full [Prescribing Information](#), including [Medication Guide](#), for BLUJEPÀ.

BLUJEPÀ 
(gepotidacin) 750 mg tablets

Have another UTI?* You're not alone

If you've ever had 2 UTIs in 6 months or 3 in a year, then you may have had recurrent UTIs.



*Here, UTI (urinary tract infection) refers to uncomplicated UTI, also known as an infection of the bladder.


Within 1 month after taking the first antibiotic

UP TO 1 IN 6 WOMEN WILL NEED
a 2nd antibiotic

or a visit to a healthcare provider
to completely resolve their UTI

Some UTI bacteria can develop
resistance to antibiotics.

This means some antibiotics may not be as
effective at treating an infection anymore.




**Does any of this
sound familiar?**

Work with your healthcare provider
to find the right UTI treatment for you

What is BLUJEPAN?

A first-of-its-kind antibiotic for uncomplicated UTIs

BLUJEPAN targets 2 bacterial enzymes in a unique way. This stops bacteria from multiplying and kills them.



Important Safety Information (cont'd)

Before starting BLUJEPAN, tell your healthcare provider if you:

- or anyone in your family has ever had a heart rhythm problem (problem with heart rate or heartbeat) or QTc prolongation.
- are taking any medicines to treat heart rhythm problems. Using these medicines with BLUJEPAN may cause serious side effects. If you are not sure, ask your healthcare provider.
- are taking medicines called moderate or strong CYP3A4 inhibitors. Using these medicines with BLUJEPAN may cause serious side effects. If you are not sure, ask your healthcare provider.

Tell your healthcare provider right away if you have any of the following symptoms:

- fast, pounding, or uneven heartbeat
- feeling dizzy
- you lose consciousness (pass out)
- feeling lightheaded
- feeling faint

• an increase of a certain chemical in your nervous system (cholinergic effects). BLUJEPAN can cause an increase of a certain chemical in your nervous system, which may cause the following symptoms:

- diarrhea
- vomiting
- fainting
- trouble saying words clearly
- muscle spasms
- nausea
- more saliva in the mouth
- more sweating
- stomach pain

Results with BLUJEPAN

In 2 clinical trials in patients with uncomplicated UTI, BLUJEPAN was effective at eliminating UTI bacteria and resolving symptoms

BLUJEPAN was compared to a proven treatment, nitrofurantoin. To show that BLUJEPAN works, the studies checked for 2 things—both had to happen, without needing another antibiotic:

- ▶ Complete resolution of UTI symptoms
- ▶ Elimination of infection-causing bacteria in the urine

Patients taking BLUJEPAN had their symptoms and bacteria eliminated similar to those taking nitrofurantoin 10 to 13 days after starting treatment:

	BLUJEPAN	Nitrofurantoin
Trial 1	52%	47%
Trial 2	59%	44%

Your results may vary.

Additional Information

Response to BLUJEPAN was also seen in patients with recurrent UTIs and against some resistant strains of UTI bacteria.

Based on limits of the analysis, results are descriptive.

Important Safety Information (cont'd)

Call your healthcare provider right away if you have trouble saying words clearly, have shortness of breath, or if you faint. These symptoms may be cholinergic effects of BLUJEPAN.

The increased chemical in your nervous system can also be connected to other symptoms. Call your healthcare provider right away if you have any of these additional symptoms:

- irregular heartbeat (heart block)
- slow heartbeat (bradycardia)
- seizures
- fainting
- chest tightness causing difficulty breathing

Please see Important Safety Information throughout and on pages 6 and 7.

Please see the accompanying full Prescribing Information, including Medication Guide, for BLUJEPAN.

BLUJEPAN
(gepotidacin) 750 mg tablets

Important Safety Information (cont'd)

• **allergic reactions.** Allergic reactions can happen in people who take BLUJEPA, including a severe allergic reaction called anaphylaxis. Stop taking BLUJEPA and get medical help right away if you have any symptoms of a severe allergic reaction, including:

- hives
- rash
- feeling faint
- swelling of your lips, tongue, or throat
- or dizzy

• **diarrhea.** Diarrhea is a common side effect caused by antibiotics including BLUJEPA. The diarrhea may happen at the beginning of your treatment with BLUJEPA and usually stops after the antibiotic is stopped. In most cases, the diarrhea will go away without treatment. In some cases, diarrhea may be caused by *Clostridiooides difficile* infection (CDI). CDI is a severe infection of the intestines (bowels) that can happen up to 2 months after finishing treatment with antibiotic medicines, including BLUJEPA. CDI can be life-threatening and can lead to death. Do not take medicines to treat diarrhea without first checking with your healthcare provider. Call your healthcare provider right away if you have the following symptoms:

- stomach cramps
- diarrhea that does not go away
- fever
- watery diarrhea
- bloody stools

Who should not take BLUJEPA?

Do not take BLUJEPA if:

you have had a severe allergic reaction to BLUJEPA.

Before taking BLUJEPA, tell your healthcare provider about all of your medical conditions, including if you:

- have heart rhythm problems. See "What is the most important information I should know about BLUJEPA?"
- have a slow heartbeat (bradycardia)
- have recently had a heart attack
- have low potassium in your blood (hypokalemia)
- are planning to have a medical or surgical procedure that puts you to sleep (general anesthesia) that will use a type of medicine given through your veins (IV) that relaxes your muscles
- have liver problems
- have kidney problems, kidney failure, or are on dialysis
- are pregnant or plan to become pregnant. It is not known if BLUJEPA will harm your unborn baby

Tell your healthcare provider if you become pregnant or think you are pregnant during your treatment with BLUJEPA.

- **Pregnancy Exposure Registry:** There is a pregnancy exposure registry for those who are exposed to BLUJEPA during pregnancy. Talk to your healthcare provider about registering with GlaxoSmithKline. You can register by calling 1-888-825-5249
- are breastfeeding or plan to breastfeed. It is not known if BLUJEPA passes into breast milk. Talk to your healthcare provider about the best way to feed your baby during your treatment with BLUJEPA

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. BLUJEPA and other medicines may affect each other, causing side effects.

Especially tell your healthcare provider if you take medicines to treat heart rhythm problems. Ask your healthcare provider or pharmacist if you are not sure if you are taking any of these medicines. Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

What are the possible side effects of BLUJEPA?

BLUJEPA can cause serious side effects, including:

- See the "What is the most important information I should know about BLUJEPA?"

The most common side effects of BLUJEPA include:

<ul style="list-style-type: none">- diarrhea- nausea- stomach pain- gas- headache	<ul style="list-style-type: none">- soft stools- dizziness- vomiting- yeast infections
---	---

These are not all the possible side effects of BLUJEPA. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report negative side effects to GSK at gsk.public.reportum.com or 1-888-825-5249.

Please see Important Safety Information throughout.

Please see the accompanying full Prescribing Information, including Medication Guide, for BLUJEPA.

BLUJEPA
(gepotidacin) 750 mg
tablets

Ask your healthcare provider about BLUJEPAP today*



Scan the QR code or visit BLUJEPAP.com to learn more

*Available by prescription only.

Important Safety Information

What is the most important information I should know about BLUJEPAP?

BLUJEPAP, an antibiotic, can cause serious side effects, including:

- **changes in electrical activity of your heart called QTc prolongation.** BLUJEPAP can cause QTc prolongation (also called prolonged QTc interval) that can lead to serious heart rhythm problems, including Torsades de Pointes. There may be changes in your heartbeat (a fast or irregular heartbeat).

Please see Important Safety Information throughout and on pages 6 and 7.

Please see the accompanying full Prescribing Information, including Medication Guide, for BLUJEPAP.

GSK for You

If you don't have prescription coverage, need cost information, or need help paying for your medicine, visit GSKforyou.com or call **1-866-GSK-FOR-U (1-866-475-3678)**.

BLUJEPAP  **GSK**
(gepotidacin)

Trademarks are owned by or licensed to the GSK group of companies.

©2026 GSK or licensor.
PMUS-GPTBROC250005 January 2026
Produced in USA.